



ARL-TR-7652 • APR 2016



# Space–Time Environmental Image Information for Scene Understanding

by Arnold Tunick

Approved for public release; distribution unlimited.

## **NOTICES**

### **Disclaimers**

The findings in this report are not to be construed as an official Department of the Army position unless so designated by other authorized documents.

Citation of manufacturer's or trade names does not constitute an official endorsement or approval of the use thereof.

Destroy this report when it is no longer needed. Do not return it to the originator.



# Space–Time Environmental Image Information for Scene Understanding

by Arnold Tunick

*Computational and Information Sciences Directorate, ARL*

REPORT DOCUMENTATION PAGE				Form Approved OMB No. 0704-0188	
<p>Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing the burden, to Department of Defense, Washington Headquarters Services, Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.</p> <p><b>PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.</b></p>					
1. REPORT DATE (DD-MM-YYYY) April 2016		2. REPORT TYPE Final		3. DATES COVERED (From - To) 08/2015–12/2015	
4. TITLE AND SUBTITLE Space–Time Environmental Image Information for Scene Understanding				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S) Arnold Tunick				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) US Army Research Laboratory ATTN: RDRL-CII-A 2800 Powder Mill Road Adelphi, MD 20783-1138				8. PERFORMING ORGANIZATION REPORT NUMBER  ARL-TR-7652	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited.					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT <p>The interpretation of space- and time-varying environmental image information can pose serious challenges for rapid and robust scene understanding, particularly for problems of interest to the Army. Changing environmental conditions, such as illumination, precipitation, and vegetation, can obscure features, degrade object recognition, and modify saliency and image context. Current methods for scene understanding do not address elements of image information (and image context) affected by space- and time-changing environmental conditions, and as a result, important and meaningful features of the image data may be overlooked. In this report, we propose that it is important to incorporate space- and time-varying environmental image information from the very beginning of the data collection process so that the recorded images can be more effectively indexed and retrieved for operational use and analysis. This top-down approach not only provides a systematic characterization of the measured data for better scene description, but can also help the end user (Soldier) develop improved course of action strategies based on scene understanding (algorithms and analysis) incorporating battlefield environments changing in space and time.</p>					
15. SUBJECT TERMS computer vision, image resolution, context, saliency, visibility, illumination, vegetation, visual motion					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT  UU	18. NUMBER OF PAGES  26	19a. NAME OF RESPONSIBLE PERSON Arnold Tunick
a. REPORT Unclassified	b. ABSTRACT Unclassified	c. THIS PAGE Unclassified			19b. TELEPHONE NUMBER (Include area code) (301) 394-1233

## Contents

---

<b>List of Figures</b>	<b>iv</b>
<b>List of Tables</b>	<b>iv</b>
<b>Acknowledgments</b>	<b>v</b>
<b>1. Introduction</b>	<b>1</b>
<b>2. Space and Time Scales</b>	<b>2</b>
<b>3. Image Resolution, Image Context, and Identifying Objects</b>	<b>3</b>
<b>4. Time- and Space-Varying Elements of Scene Understanding</b>	<b>6</b>
4.1 Environmental Image Information	7
4.2 Scene Description Indexing	8
<b>5. Summary and Conclusions</b>	<b>10</b>
<b>6. References</b>	<b>11</b>
<b>List of Symbols, Abbreviations, and Acronyms</b>	<b>17</b>
<b>Distribution List</b>	<b>18</b>

## List of Figures

---

Fig. 1	Primary space ( $s$ ) and time ( $t$ ) scales .....	3
Fig. 2	Can you correctly identify these images? Image resolution: a) 30 x 20 pixels, b) 30 x 14 pixels, and c) 30 x 16 pixels.....	4
Fig. 3	Higher-resolution image scenes corresponding to the 3 shapes shown in Fig. 2. a) Taj Mahal (photo courtesy of desktopdress.com), b) US Capitol Dome (photo courtesy of Library of Congress), and c) nuclear power plant, Bushehr, Iran (photo courtesy of Behrouz Mehri/AFP/Getty Images). .....	4
Fig. 4	Can you correctly identify these objects? .....	5
Fig. 5	Low-resolution images of the scene from which the objects in Fig. 4 were taken, where neither large nor small objects are discernible (left: 16 x 10 image pixels, right: 32 x 20 image pixels) .....	5
Fig. 6	Same images as shown in Fig. 5 but with a slightly higher resolution (left: 64 x 40 image pixels, right: 128 x 80 image pixels) .....	5
Fig. 7	Same images as shown in Figs. 5 and 6 but with an even higher resolution (left: 525 x 336 image pixel, right: 3888 x 2492 image pixels). Note that the hoist, surveyor, and engineers wearing hard hats in the far-field of the imaged scene all required increased image resolution to be clearly identified. ....	6

## List of Tables

---

Table 1	Image resolution information (in numbers of pixels) .....	6
Table 2	Time- and space-varying elements of scene understanding .....	8
Table 3	Scene description indexing (top-level view).....	9
Table 4	Available/accessible environmental image information .....	9

## **Acknowledgments**

---

I thank RE Meyers, P David, and G Warnell for helpful discussions. This research was supported by the US Army Research Laboratory.

INTENTIONALLY LEFT BLANK.



## 1. Introduction

---

Rapid and robust scene understanding is a critically important goal for the development of Army autonomous intelligent systems.<sup>1</sup> For outdoor natural scenes, it will be important for autonomous intelligent systems to be able to quickly discern the depth of view, navigability, exposure or concealment (as it relates to object searching), and transience, that is, the rate at which elements of the scene or its environment are changing in space and time.<sup>2,3</sup> In this regard, saliency estimation has been helpful to computationally identify elements in a scene that immediately capture the visual attention of an observer.<sup>4,5</sup> Several recent papers have discussed concepts associated with visual saliency to enhance automated navigation and scene exploration.<sup>6-8</sup> Note, however, that the most active or salient object(s) in a scene may not represent the most important or meaningful feature(s) of the scene.<sup>9</sup> For example, an automated vision system may readily detect changes in the ground surface as a new or different object in the field of view; however, recognizing the physical characteristics of the new surface<sup>10,11</sup> (e.g., shallow or deep water, thick or thin ice, snow, mud, quicksand, etc.) and observing any changes in the environmental context of the image<sup>12-14</sup> may be critically important. Characterizing interactions between objects and the environment also can contribute to physical scene understanding.<sup>15,16</sup> In the example above, if a scene depicts vehicles or personnel activity in a changing complex environment, then robust scene understanding could provide key border and accessibility information for navigation and trafficability.

Nevertheless, many current methods for scene understanding, like those that generate image descriptions via automated semantic labeling<sup>17</sup> or visual scene classification,<sup>18</sup> do not address image information (and image context) affected by changing environmental conditions. Yet, the interpretation of changing environmental conditions can pose serious challenges for computer vision processes, such as those associated with place recognition, navigation, road/terrain detection, and scene exploration.<sup>19-24</sup> This is because rain, snow, and fog weather events, smoke, haze, or other changes in lighting and visibility can significantly obscure features, degrade object recognition, and modify the saliency and image context of an outdoor scene.<sup>25-32</sup> Naturally, scene-depicted environmental conditions can vary with time of day, season, and location.<sup>33</sup>

Similar challenges can also extend to interpreting space- and time-changing scenes due to visual motion of objects within the field of view.<sup>34-36</sup> In this case, changing environmental conditions such as illumination, precipitation, and vegetation can make feature recognition of moving objects unclear, so that identifying moving objects in outdoor environments becomes more difficult for vision-based intelligent

systems.<sup>35,36</sup> As an example, poor contrast in images can be brought about by low visibility due to environmental effects or weak illumination, such as during dawn, dusk, or night.<sup>34</sup> Visually degraded or blurred images can be brought about by rapid movements of the camera and/or objects in the field of view, especially in the low-light case, which necessitate longer camera exposure times.<sup>21,34,37</sup> In addition, space and time variations in scene illumination can affect the optical flow field in images and movies.<sup>38</sup> Note that there have been many optical flow approaches used to detect the motion of objects in a scene, which have been helpful in a variety of applications.<sup>39-41</sup> Nevertheless, camera motion may introduce some unmanageable artifacts with some of these gradient-based optical flow approaches if they are not augmented by more sophisticated spatiotemporal analyses.<sup>42-44</sup>

In this report, we propose that it is important to incorporate space- and time-varying environmental image information from the very beginning of the data collection process so that the recorded images can be more effectively indexed and retrieved for operational use and analysis. This top-down approach not only provides a systematic characterization of the measured data for better scene description, but can help the end user (Soldier) develop improved course of action strategies based on scene understanding (algorithms and analysis) incorporating battlefield environments changing in space and time. Incorporating space- and time-varying environmental image information for better scene understanding can be vital to support numerous Army missions,<sup>45-49</sup> such as those related to weather elements on the battlefield that can alter terrain features and trafficability; low visibility that can impede reconnaissance and target acquisition or alternately conceal friendly forces maneuvers and activities; and wind speed and direction that can favor upwind forces in nuclear, biological, and chemical (NBC) attacks or decrease the effectiveness of downwind forces due blowing dust, smoke, sand, rain, or snow. In addition, reporting wind speed and direction information at the time images are being recorded can significantly influence the success of aviation-related missions, like those associated with unmanned aerial vehicle take off, landing, and in-flight control.<sup>50</sup>

## **2. Space and Time Scales**

---

Based on an analysis by Meyers,<sup>12</sup> this section provides a framework to help categorize the spatial and temporal properties of recorded image data. Relevant time scales include, but are not limited to, the shutter exposure time, time interval between frames, time over which images are captured in a sequence, and time over which there is visual motion of objects inside the field of view. Space scales include, but are not limited to, the field of view, depth of view, image resolution, pixel size, pixel separation, scene color or shading variations as a function of spatial

location, spatial smearing of moving elements in the field of view, spatial smearing due to optical turbulence and environmental/weather effects, and smearing of textures in the field of view. Naturally, the smearing of elements in the field of view can also be related to the temporal resolution of the image data. Figure 1 illustrates the primary space and time scales that can be used to describe the various spatial and temporal resolutions of objects and/or activities in a recorded image scene, to include images recorded in varying environmental conditions. Here,  $\Delta s$  and  $\Delta t$  represent changes in position and time, respectively.





Fig. 1 Primary space ( $s$ ) and time ( $t$ ) scales


### 3. Image Resolution, Image Context, and Identifying Objects

To start, let us explore the impact of varying image resolution and image context on the analysis of an outdoor scene. Try to identify the 3 dome shapes in Fig. 2. Without some additional information related to the object size, texture, or shape or knowledge of how the image context may change in an expanded field of view, it is difficult to correctly identify and label these familiar images. Furthermore, distinguishing various image details, even in ideal conditions with regard to lighting and visibility, can depend on the image contrast and resolution, where image resolution here refers to the number of pixels that compose the image data input. Interestingly, Torralba<sup>51</sup> reported that for human vision the brain can comprehend the gist of an image scene remarkably quickly, regardless of whether low- or high-resolution images are used. He concluded that images at the resolution of 32 x 32 color pixels can provide an observer enough information to correctly identify the semantic category and general layout of an indoor/outdoor scene. For example, in

Fig. 2 the main “dome” category for these low-resolution images is identifiable. However, if we consider Fig. 3, which contains expanded fields of view and higher resolution images from which the elements in Fig. 2 were taken, then the building domes and many additional image details can be identified over a much wider range of spatial scales.



**Fig. 2** Can you correctly identify these images? Image resolution: a) 30 x 20 pixels, b) 30 x 14 pixels, and c) 30 x 16 pixels.




**Fig. 3** Higher-resolution image scenes corresponding to the 3 shapes shown in Fig. 2. a) Taj Mahal (photo courtesy of desktopdress.com), b) US Capitol Dome (photo courtesy of Library of Congress), and c) nuclear power plant, Bushehr, Iran (photo courtesy of Behrouz Mehri/AFP/Getty Images).

To demonstrate this point further, Can you identify the 2 similarly shaped objects shown in Fig. 4 without some additional context? What if we look at the complete image (Fig. 5) from which the objects were taken? In this case, at low resolution, it is quite difficult to discern any individual elements in the field of view. Of course, the degree of image resolution needed for a particular task depends on the analysis or computer vision problem of interest.<sup>5,8,17,18,21,52–55</sup> Yet, with regard to scene understanding and semantic labeling, the slightly higher resolution images shown in Fig. 6 clearly provide more usable information. In other words, when the image resolution is increased to 64 x 40 pixels and greater, one can more easily identify the layout and main elements of the image scene, such as the reactor dome and hard hat shown above. However, if still higher resolution images of this reactor site are considered (Fig. 7), then additional details and information may be gained, for example, intelligence relating to its operational status. By analyzing the extracted and labeled objects shown in Fig. 7, one might ask if the reactor site is still under


construction or near completion as evidenced by the engineers wearing hard hats, the surveyor, the hoist, and the electrical hazard sign. Note here that the hoist, surveyor, and engineers wearing hard hats in the far-field of the imaged scene all required increased resolution (i.e.,  $\geq 32 \times 32$  pixels) to be clearly identified (visually compare right vs. left in Fig. 7). Table 1 provides the various image resolution details (in numbers of pixels) for the labeled objects in this outdoor scene.




**Fig. 4** Can you correctly identify these objects?



**Fig. 5** Low-resolution images of the scene from which the objects in Fig. 4 were taken, where neither large nor small objects are discernible (left: 16 x 10 image pixels, right: 32 x 20 image pixels)



**Fig. 6** Same images as shown in Fig. 5 but with a slightly higher resolution (left: 64 x 40 image pixels, right: 128 x 80 image pixels)



**Fig. 7** Same images as shown in Figs. 5 and 6 but with an even higher resolution (left: 525 x 336 image pixel, right: 3888 x 2492 image pixels). Note that the hoist, surveyor, and engineers wearing hard hats in the far-field of the imaged scene all required increased image resolution to be clearly identified.

**Table 1** Image resolution information (in numbers of pixels)

Object	Fig. 7 (left)	Fig. 7 (right)
Main image	525 x 336	3888 x 2492
Reactor dome	191 x 51	1028 x 256
Hard hat	82 x 45	405 x 225
Danger sign	32 x 36	64 x 69
Hoist	5 x 24	39 x 175
Surveyor	5 x 11	34 x 74
Engineers	5 x 5	32 x 41

Thus, we have visually demonstrated that image resolution and image context play an important role in being able to clearly recognize and identify individual objects for scene understanding. Here, object recognition may be achieved through applications associated with deep learning neural networks.<sup>56–58</sup> Also, it is important to note that image context information can include elements related to time, geographical location, and environmental conditions,<sup>12–14</sup> which can help to provide a more detailed description of recorded scenes for indexing and future retrieval, as is discussed next.

#### **4. Time- and Space-Varying Elements of Scene Understanding**

While image resolution is a key element for identifying objects and quickly discerning the gist and general layout of a recorded scene, there are many other time- and space-varying elements that are equally important for scene understanding that should be addressed from the very beginning of the data collection process.

## 4.1 Environmental Image Information

---

There are many key pieces of information that can be identified as new image data are being recorded that are important and accessible, but are usually overlooked or left undocumented. For example, one can readily identify a timestamp (relative to the sun's angle or relative to a world clock), the global positioning system (GPS) position, the prevailing environmental and weather conditions, and the field of view, depth of view, and image resolution (Table 2). The first group, shown in Table 2, focuses on environmental information, such as the GPS position and altitude above ground level (AGL), prevailing weather, cloud cover, ground and road conditions, and visibility (e.g., fog, smoke, haze, obscurants, or optical turbulence).

Identifying the key environmental and terrain conditions can provide location and geographical context information to help categorize image scenes recorded in diverse regions (e.g., coastal, mountain-valley, desert, forest, urban, rural, ocean, and arctic). Detailed terrain characteristics (e.g., muddy, sandy, gravelly, wet, dry, or icy) and reports of the most current weather conditions available (e.g., rain, snow, fog, or haze) can be retrieved and annotated to help describe images that may be used to support the planning and/or execution of military operations, as mentioned above. Typically, changing weather conditions, cloud cover, and visibility will bring about changes in the illumination of a scene, which can affect image contrast and resolution.<sup>25,30,31</sup> Time of day and sun angle information also can be retrieved, which can be useful to indicate when glare, shadows, or silhouettes may cause difficulties for computer vision processes.<sup>52,59,60</sup> Also, taking note of optical turbulence conditions is important because these effects can significantly degrade and blur image quality due to spatial smearing.<sup>61</sup>

The second group in Table 2 lists elements related to the camera specifications and the image data measurements themselves (e.g., the spatial and temporal image resolutions, field of view, depth of view, and scene color or shading variations). Together with the environmental information, these elements can be used as a basic building block for detailed scene description and image indexing.



**Table 2 Time- and space-varying elements of scene understanding**

<b>Environmental information</b>
GPS position and altitude AGL
Location: geographical context
Timestamp
Weather conditions, sky and cloud cover
Sun/moon angle
Ground/road conditions
Visibility
Vegetation
Buildings, parking lots, people, or crowds
<b>Image/camera information</b>
Image resolution
Pixel size and pixel separation
Scene color or shading variations
Field of view and depth of view
Shutter exposure time
Time interval between image frames
Time over which images are captured in a sequence

## 4.2 Scene Description Indexing

Based on the time- and space-varying elements for scene understanding described above, Table 3 provides a top-level view for scene description indexing (i.e., these are the questions that one should endeavor to address as image data are being recorded so that the data can be best indexed and retrieved for later use). In most cases, the image information (right column) can be annotated based on the camera type, lensing, pixel array, and timing specifications. Also, for example, co-located range finder instrumentation could provide effective depth and field of view measurements for this purpose. Additionally, when communications are available, the most current environmental information available (left column) can be extracted from several accessible resources, such as those shown in Table 4. Obtaining the most current data available is advantageous since environmental conditions (e.g., weather and terrain) can change over very short temporal and spatial intervals. For example, access to data from the Department of Defense (DOD) GPS<sup>62</sup> can provide latitude and longitude or Universal Transverse Mercator (UTM) location and timestamp information, commonly reported as Greenwich Mean Time (GMT) or Coordinated Universal Time (UTC). Also, data from the US Naval Observatory (USNO)<sup>63</sup> can provide precise timing information as well as solar and lunar elevation/azimuth angles. Similarly, terrain and geographical location and context information can be provided by satellite and aerial imagery from the US Army Corps of Engineers, Army Geospatial Center (USACE AGC)<sup>64</sup> for military operations or from public Internet resources such as Google,<sup>65</sup> MapQuest,<sup>66</sup> Bing,<sup>67</sup> and Yahoo Maps.<sup>68</sup>



**Table 3     Scene description indexing (top-level view)**

	<b>Environmental information</b>	<b>Image/camera information</b>
1	What is the GPS position of the depicted scene?	What is the camera field of view?
2	What is the altitude (AGL) of the depicted scene?	What is the scene depth of view?
3	What is the timestamp of the recorded image?	What is the image spatial resolution (in pixels)?
4	What are the current weather conditions?	What is the camera pixel size?
5	What is the percent sky/cloud cover?	What is the scene color (R, B, G or grayscale)?
6	What is the sun elevation/azimuth angle?	What are the spatial shading variations of the scene?
7	What is the visibility?	What is the camera integration time?
8	What are the current ground/road conditions?	What is the camera shutter exposure time?
9	What vegetation is in the field of view?	What is the time interval between frames?
10	What buildings, parking lots, people or crowds are in the field of view?	What is the starting, ending and total time for the recorded image sequence?

**Table 4     Available/accessible environmental image information**

1	DOD GPS: Latitude/longitude or UTM, altitude (AGL), GMT, or UTC
2	USNO: Precise time, sun/moon elevation/azimuth angle
3	Terrain and location: USACE AGC — Satellite/aerial imagery and terrain analysis
4	Terrain and location: Google, MapQuest, Bing, Yahoo Maps
5	Weather: USAF 557th Weather Wing
6	Weather: National Weather Service (NWS) and National Centers for Environmental Information (NCEI)
7	Weather: Intellicast, AccuWeather, Weather Underground

Weather conditions and related oceanic, atmospheric, and geophysical data are also available for the military through the US Air Force 557th Weather Wing<sup>69</sup> (i.e., formerly the US Air Force Weather Agency) and for the civilian community through the NWS<sup>70</sup> and NCEI.<sup>71</sup> Daily NWS weather reports can be found online containing hourly records citing the date, time, wind speed (miles per hour), visibility (miles), weather (i.e., rain, snow, fog, haze, etc.), sky/cloud condition (reported as overcast [OVC], broken [BRK], scattered [SCT], or clear [CLR] along with the cloud ceiling height in hundreds of feet AGL), air temperature, dew point temperature, relative humidity (%), pressure, and precipitation (in inches). Naturally, current weather and weather forecast information are readily found on Internet web sites, such as Intellicast,<sup>72</sup> AccuWeather,<sup>73</sup> and Weather Underground.<sup>74</sup> Note however, that in areas where communications are either restricted or unavailable, the information needed to describe the scene, i.e., as outlined in Table 3, should instead be gleaned from the recorded images when they are retrieved for analysis.

Thus, we have shown that abundant time- and space-varying environmental image information can be accessed and annotated to augment image data as they are being recorded for a better organized, top-down approach to scene description, indexing and image retrieval.

## 5. Summary and Conclusions

---

In this report, we proposed that it is important to incorporate space- and time-varying environmental image information at the start of the data collection process in order to provide the end user (Soldier) with a better organized, top-down approach to index and retrieve image data for operational use and analysis. We provided several examples to show that space- and time-varying elements of environmental image information (and changes in image context) can be used as a basic building block for detailed scene description, and as such, could be used to support Army mission planning and execution. In conclusion, we anticipate that incorporating space- and time-varying environmental image information in the data measurement process will lead to 1) improved autonomous intelligent systems supporting Army missions in complex and changing environments and 2) improved course of action strategies based on scene understanding (algorithms and analysis) incorporating battlefield environments changing in space and time.

## 6. References

---

1. Army Research Laboratory (US). US Army Research Laboratory S&T Campaign Plans 2015–2035, Adelphi (MD): Army Research Laboratory (US); 2014. System Intelligence & Intelligent Systems; p. 98–99.
2. Greene MR, Oliva A. Recognition of natural scenes from global properties: seeing the forest without representing the trees. *Cog Psych.* 2009;58:137–176.
3. Greene MR, Oliva A. The briefest of glances: the time course of natural scene understanding. *Psych Sci.* 2009;20(4):464–472.
4. Itti L, Koch C, Niebur E. A model of saliency-based visual attention for rapid scene analysis. *IEEE Trans Pattern Analysis and Machine Intel.* 1998;20(11):1254–1259.
5. Perazzi F, Krahenbuhl P, Pritch Y, Hornung A. Saliency filters: contrast based filtering for salient region detection. *Proceedings of IEEE Conference on Computer Vision and Pattern Recognition*; 2012 Jun 16–21; Providence, RI.
6. Roberts R, Ta D-N, Straub J, Ok K, Dellaert F. Saliency detection and model-based tracking: a two part vision system for small robot navigation in forested environment. *Proc. SPIE 8387, Unmanned Systems Technology XIV Conference*; 2012 May 1. Bellingham (WA): The International Society for Optical Engineering; c2012.
7. Yeomans B, Shaukar A, Gao Y. Testing saliency based techniques for planetary surface scene analysis. In: *ASTRA 2015. Proceedings of 13th Symposium on Advanced Space Technologies in Robotics and Automation*; 2015 May 11–13; Noordwijk (The Netherlands).
8. Warnell G, David P, Chellappa R. Ray saliency: bottom-up visual saliency for a rotating and zooming camera. *Intl J Computer Vis.* 2015. doi 10.1007/s11263-015-0842-9.
9. Hung HSW. From visual saliency to video behaviour understanding [doctoral dissertation]. [London (UK)]: Queen Mary, University of London; 2007.
10. Adelson EH. On seeing stuff: the perception of materials by humans and machines. *Proc. SPIE 4299, Conference on Human Vision and Electronic Imaging VI Conference*; 2001 Jan 20; San Jose (CA). Bellingham (WA): The International Society for Optical Engineering; c2001.
11. Siva P, Russell C, Xiang T, Agapito L. Looking beyond the image: Unsupervised learning for object saliency and detection. In: *CVPR 2013*.

- Proceedings of IEEE Conference on Computer Vision and Pattern Recognition; 2013 Jun 23–28; Portland (OR).
12. Meyers RE. Army Research Laboratory (US), Adelphi, MD. Personal communication, 2014.
  13. Rawat YS, Kankanhalli MS. Context-aware photography learning for smart mobile devices. *ACM Trans Multimedia Computing, Comms, and Applications* 2105 Oct;12(1s): Article 19.
  14. Matas J, Murino V, Leal-Taixe L, Rosenhahn B., editors. Holistic scene understanding. Report from the Dagstuhl Seminar 15081, 2015 Feb 15–20; Wadern, Germany.
  15. Battaglia PW, Hamrick JB, Tenenbaum JB. Simulation as an engine of physical scene understanding. *Proc National Academy of Sciences*. 2013;110(45):18327–18332.
  16. Ullman TD, Stuhlmüller A, Goodman ND, Tenenbaum JB. Learning physics from dynamical scenes. In: *CSS 2014. Proceedings of 36th Annual Conference of the Cognitive Science Society*. 2014 Jul 23–26; Quebec City, Canada.
  17. Karpathy A, Fei-Fei L. Deep visual-semantic alignments for generating image descriptions. *arXiv:1412.2306v2*; 2015.
  18. Wigness M, Draper BA, Beveridge JR. Efficient label collection for unlabeled image datasets. In: *CVPR 2015. Proceedings of IEEE Conference on Computer Vision and Pattern Recognition*; 2015; Boston, MA. p. 4594–4602.
  19. Katsura H, Miura J, Hild M, Shirai Y. A view-based outdoor navigation using object recognition robust to changes of weather and seasons. In: *IROS 2003. Proceedings of IEEE International Conference on Intelligent Robots and Systems*; 2003; Las Vegas, NV. p. 2974–2979.
  20. Milford MJ, Wyeth GF. SeqSLAM: Visual route-based navigation for sunny summer days and stormy winter nights. In: *ICRA 2012. Proceedings of IEEE International Conference on Robotics and Automation*. 2012; Saint Paul, MN.
  21. Milford M. Vision-based place recognition: how low can you go? *Intl J Robotics Res*. 2013;32(7):766–789.
  22. Milford M, Vig E, Scheirer W, Cox D. Vision-based simultaneous localization and mapping in changing outdoor environments. *J Field Robotics*. 2014;31(5):780–802.

23. Alvarez JM, Lopez AM, Gevers T, Lumberras F. Combining priors, appearance and context for road detection. *IEEE Trans Intelligent Transportation Sys.* 2014;5(3):1168–1178.
24. Scharwachter T, Franke U. Low-level fusion of color, texture and depth for robust road scene understanding. *IEEE Intelligent Vehicles Symposium IV*; 2015 Jun 28–Jul 1; Seoul, Korea.
25. Narasimhan SG, Nayar SK. Vision and the atmosphere. *Intl J Computer Vision.* 2002;48(3):233–254.
26. Narasimhan SG, Wang C, Nayar SK. All the images of an outdoor scene. In: *Proceedings of 7th European Conference on Computer Vision*; 2002; Copenhagen, Denmark. p. 148–162.
27. Lalonde J-F, Efros AA, Narasimhan SG. Estimating the natural illumination conditions from a single outdoor image. *Intl J Computer Vis.* 2012;98:123–145.
28. Xie L. Geographic and environmental interpretation of photographs [master's thesis]. Merced (CA): University of California, Merced; 2011.
29. Yao C, Wang C, Hong L, Cheng Y. A Bayesian probabilistic framework for rain detection. *Entropy.* 2014;16:3302-3314.
30. He K, Sun J, Tang X. Single image haze removal using dark channel prior. *IEEE Trans Pattern Analysis and Machine Intel.* 2011;33(12):2341–2353.
31. Song Y, Luo H, Hui B, Chang Z. An improved image dehazing and enhancing method using dark channel prior. In: *Proceedings of 27th Chinese Control and Decision Conference.* 2015 May 23–25; Qingdao, China. p. 5840 – 5845.
32. Lu C, Lin D, Jia J, Tang C-K. Two-class weather classification. In: *CVPR 2014. Proceedings of IEEE Conference on Computer Vision and Pattern Recognition.* 2014; Columbus, OH. p. 3718–3725.
33. Yu J, Luo J. Leveraging probabilistic season and location context models for scene understanding. In: *Proceedings of the International Conference on Content-Based Image and Video Retrieval.* 2008; Niagara Falls, Ontario (Canada). p. 169–178.
34. Teutsch M. Moving object detection and segmentation for remote aerial video surveillance [doctoral dissertation]. Karlsruhe (Germany): Karlsruhe Institute of Technology; 2014.

35. Holtzhausen PJ, Crnojevic V, Berbst BM. An illumination invariant framework for real-time foreground detection. *J Real-Time Image Process.* 2015;10:423–433.
36. Tu Z, Zheng A, Yang E, Luo B, Hussain A. A biologically inspired vision-based approach for detecting multiple moving objects in complex outdoor scenes. *Cog Computation.* 2015. doi 10.1007/s12559-015-9318-z.
37. Kim MD, Ueda J. Dynamics-based motion de-blurring for a PZT-driven, compliant camera orientation mechanism. *Intl J Robotics Res.* 2015;34:653–673.
38. Andreev R, Scherzera O, Zulehner W. Simultaneous optical flow and source estimation: Space—time discretization and preconditioning. *Applied Numerical Math.* 2015;96:72–81.
39. Fortun D, Bouthemy P, Kervrann C. Optical flow modeling and computation: A survey. *Computer Vis and Image Understanding.* 2015;134:1–21.
40. Fleet DJ, Weiss Y. *Mathematical Models in Computer Vision: The Handbook.* Berlin (Germany): Springer; 2005; p. 239–258. Chapter 15, Optical flow estimation.
41. Barron J, Fleet D, Beauchemin S. Performance of optical flow techniques. *Intl J Computer Vis.* 2004;12(1):43–77.
42. Laptev I. Local spatio-temporal image features for motion interpretation [doctoral dissertation]. Stockholm (Sweden): KTH Royal Institute of Technology; 2004. isbn: 91-7283-793-4.
43. Fagerstrom D. Galilean differential geometry of moving images. In: *Computer Vision – ECCV 2004*; 2004. Berlin, Heidelberg (Germany): Springer-Verlag p. 494–506.
44. Fagerstrom D. Spatio-temporal scale-space theory [doctoral thesis]. Stockholm (Sweden): KTH Royal Institute of Technology; 2011. isbn: 978-91-7501-024-3.
45. Headquarters, Department of the Army. Weather Support for Army Tactical Operations, Appendix B – Weather Effects on Army Operations. Washington (DC): Headquarters, Department of the Army; 1989 Aug 31. Field Manual No.: FM 34-81.
46. Headquarters, Department of the Army. Field Behavior of NBC Agents (Including Smoke and Incendiaries). Washington (DC): Headquarters, Department of the Army; 1986 Nov 3. Field Manual No.: FM 3-6.

47. Headquarters, Department of the Army. Weather Support and Services for the US Army. Washington (DC): Headquarters, Department of the Army; 2015 Sep 10. Field Manual No.: AR 115-10.
48. Datz IM. Military operations under special conditions of terrain and weather. New Delhi (India): Lancer Publishers; 2008.
49. Szymber RJ. US Army tactical weather support requirements for weather and environmental data elements and meteorological forecasts. Adelphi (MD): Army Research Laboratory (US); 2006 Feb. Report No.: ARL-TR-3720.
50. Headquarters, Department of the Army. Army Unmanned Aircraft System Operations. Washington (DC): Headquarters, Department of the Army; 2009 Jul 29. Field Manual No.: FM 3-04.155.
51. Torralba A. How many pixels make an image? *Vis Neuroscience*. 2009;26:23–131.
52. Olszewska JI. Semantic, automatic image annotation based on multi-layered active contours and decision trees. *Intl J Advanced Computer Sci and Applications*. 2013;4(8):201–208.
53. Fleuret F, Geman D. Coarse-to-fine face detection. *Intl J Computer Vis*. 2001;41:85–107.
54. Park D, Ramanan D, Fowlkes C. Multiresolution models for object detection. In: *Computer Vision - ECCV 2010*; 2010. Berlin, Heidelberg (Germany): Springer-Verlag. p. 241–254.
55. Pedersoli M, Vedaldi A, Gonzalez J. A coarse-to-fine approach for fast deformable object detection. In: *CVPR 2011. IEEE Computer Vision and Pattern Recognition Conference*; 2011 Jun 21–23; Colorado Springs, CO.
56. Bergstra J, Breuleux O, Bastien F, Lamblin P, Pascanu R, Desjardins G, Turian J, Warde-Farley D, Bengio Y. Theano: A CPU and GPU math expression compiler. In: *Proceedings of the Python for Scientific Computing Conference*; 2010 Jun 30–Jul 3; Austin, TX.
57. Bastien F, Lamblin P, Pascanu R, Bergstra J, Goodfellow I, Bergeron A, Bouchard N, Warde-Farley D, Bengio Y. Theano: new features and speed improvements. *Neural Information Processing Systems, Deep Learning Workshop*; 2012 Dec 8; Lake Tahoe, CA.
58. Ding W, Wang R, Mao F, Taylor G. Theano-based large-scale visual recognition with multiple GPUs. *arXiv:1412.2302v4*; 2015.

59. Shafer SA. Shadows and silhouettes in computer vision. Berlin (Germany): Kluwer Academic Publishers; 1985.
60. Reddy D, Veeraraghavan A. Lens flare and lens glare. In: Computer Vision: A Reference Guide; 2014. Berlin (Germany): Springer. p. 445–447.
61. Roggemann MC, Welsh BM, Hunt BR. Imaging through turbulence. Boca Raton (FL): CRC Press; 1996.
62. GPS.gov. US Air Force; 8 Dec 2015 [accessed 2015 Dec]. <http://www.gps.gov/governance/agencies/defense/>.
63. The United States Naval Observatory (USNO). US Navy [accessed 2015 Dec]. <http://www.usno.navy.mil/USNO>.
64. Army Geospatial Center. US Army Corps of Engineers [accessed 2015 Dec]. <http://www.agc.army.mil/>.
65. Google Maps. [accessed 2015 Dec]. <https://www.google.com/maps/>.
66. MapQuest. [accessed 2015 Dec]. <http://www.mapquest.com/>.
67. Bing Maps. [accessed 2015 Dec]. <https://www.bing.com/maps/>.
68. YAHOO! Maps. [accessed 2015 Dec]. <https://maps.yahoo.com/b/>.
69. 557<sup>th</sup> Weather Wing. US Air Force [accessed 2015 Dec]. <http://www.557weatherwing.af.mil/>.
70. National Weather Service [accessed 2015 Dec]. <http://www.weather.gov/>.
71. National Centers for Environmental Information (NCEI). National Oceanic and Atmospheric Administration [accessed 2015 Dec] <https://www.ncei.noaa.gov/>.
72. Intellicast. The Weather Company LLC; 2016 [accessed 2015 Dec]. <http://www.intellicast.com/>.
73. AccuWeather.com. 2016 [accessed 2015 Dec]. <http://www.accuweather.com/>.
74. Weather Underground. The Weather Company LLC; 2016 [accessed 2015 Dec]. <http://www.wunderground.com/>.



## List of Symbols, Abbreviations, and Acronyms

---

AGC	Army Geospatial Center
AGL	above ground level
BRK	broken
CLR	clear
DOD	Department of Defense
GMT	Greenwich Mean Time
GPS	global positioning system
NCEI	National Centers for Environmental Information
NWS	National Weather Service
OVC	overcast
SCT	scattered
USACE	US Army Corps of Engineers
USNO	US Naval Observatory
UTC	Coordinated Universal Time
UTM	Universal Transverse Mercator

1 DEFENSE TECHNICAL  
(PDF) INFORMATION CTR  
DTIC OCA

2 DIRECTOR  
(PDF) US ARMY RESEARCH LAB  
RDRL CIO LL  
IMAL HRA MAIL & RECORDS  
MGMT

1 GOVT PRINTG OFC  
(PDF) A MALHOTRA

7 DIRECTOR  
(PDF) US ARMY RESEARCH LAB  
RDRL CII  
B BROOME  
RDRL CII A  
S YOUNG  
A TUNICK  
G WARNELL  
P DAVID  
RDRL CIN T  
R MEYERS  
K DEACON